



**HOLMES FAMILY MEDICINE**  
 151 PARKVIEW DRIVE  
 MILLERSBURG, OHIO 44654  
 PH 330-674-1200 FAX 330-674-3320

**DRUG DOSAGES**

<b>Tylenol (Acetaminophen)</b> (dosing interval: every 4-6 hours as needed)					
			<b>Infant/Childrens Suspension 160/5ml</b>	<b>Chewables 80 mg</b>	<b>Junior Strength Chewables 160mg</b>
<b>Weight:</b>	<b>Age:</b>				
6-11 lbs	0-3 mos		1/4 tsp/1.25 ml	-	-
12-17 lbs	4-11 mos		1/2 tsp/2.5 ml	-	-
18-23 lbs	12-23 mos		3/4 tsp/3.75 ml	-	-
24-35 lbs	2-3 yrs		1 tsp/5 ml	2	-
36-47 lbs	4-5 yrs		1.5 tsp/7.5 ml	3	-
48-59 lbs	6-8 yrs		2 tsp/10 ml	4	2
60-71 lbs	9-10 yrs		2.5 tsp/12.5 ml	5	2 and 1/2
72-95 lbs	11 yrs		3 tsp/15 ml	6	3
96 lbs & over	12 yrs		-	-	4
<b>Children's Advil or Motrin (Ibuprofen)</b> (dosing interval: every 6 hours as needed)					
			<b>Infant Drops 50 mg/1.25 ml (Use measurements on dropper)</b>	<b>Children's Suspension 100 mg per 5 ml (1 teaspoon)</b>	<b>Junior Strength Advil Tablets 100 mg / tablet</b>
<b>Weight:</b>	<b>Age:</b>				
13-17 lbs	6-11 mos		1.25 ml	-	-
18-23 lbs	12-23 mos		1.875 ml	-	-
24-35 lbs	2-3 yrs		-	1 tsp (5 ml)	1 tablet
36-47 lbs	4-5 yrs		-	1 and 1/2 tsp (7.5 ml)	1.5 tablets
48-59 lbs	6-8 yrs		-	2 tsp (10 ml)	2 tablets
60-71 lbs	9-10 yrs		-	2 and 1/2 tsp (12.5 ml)	2 1/2 tablets
72-95 lbs	11 yrs		-	3 tsp (15 ml)	3 tablets

**Over-the-counter cough and cold preparations are not available for children under 6 years of age due to concerns about the side effects of the medications and a lack of benefit in improving symptoms. If you are giving a cough or cold medication to a child age 6 or older, please follow the instructions on the packaging that comes with it.**

**If your young child has cold symptoms, we recommend using saline nasal spray (with bulb syringe in young infants), Vicks VapoRub, and a cool mist humidifier or vaporizer.**

**If fever and/or symptoms have not improved in 24 hours, please contact your physician.**